arXiv:quant-ph/0306012v2 26 Nov 2003

Systems of orthogonal polynomials defined by hypergeometric type equations with application to quantum mechanics

Nicolae Cotfas

Faculty of Physics, University of Bucharest, PO Box 76-54, Postal Office 76,
Bucharest, Romania,
E-mail address: ncotfas@yahoo.com http://fpcm5.fizica.unibuc.ro/∼ncotfas

Abstract. A hypergeometric type equation satisfying certain conditions defines either a finite or an infinite system of orthogonal polynomials. We present in a unified and explicit way all these systems of orthogonal polynomials, the associated special functions and the corresponding raising/lowering operators. The considered equations are directly related to some Schrödinger type equations (Pöschl-Teller, Scarf, Morse, etc), and the defined special functions are related to the corresponding bound-state eigenfunctions.

1. Introduction

Many problems in quantum mechanics and mathematical physics lead to equations of the type

$$\sigma(s)y''(s) + \tau(s)y'(s) + \lambda y(s) = 0 \tag{1}$$

where $\sigma(s)$ and $\tau(s)$ are polynomials of at most second and first degree, respectively, and λ is a constant. These equations are usually called *equations of hypergeometric type* [5], and each of them can be reduced to the self-adjoint form

$$[\sigma(s)\varrho(s)y'(s)]' + \lambda\varrho(s)y(s) = 0 \tag{2}$$

by choosing a function ϱ such that $[\sigma(s)\varrho(s)]' = \tau(s)\varrho(s)$.

The equation (1) is usually considered on an interval (a, b), chosen such that

$$\begin{array}{rll} \sigma(s) > 0 & \text{for all} & s \in (a,b) \\ \varrho(s) > 0 & \text{for all} & s \in (a,b) \\ \lim_{s\to a}\sigma(s)\varrho(s) = \lim_{s\to b}\sigma(s)\varrho(s) = 0. & & \end{array} \tag{3}$$

Since the form of the equation (1) is invariant under an affine change of variable, it is sufficient to analyse the cases presented in table 1. Some restrictions are to be imposed to α, β in order the interval (a, b) to exist.

Table 1. The main particular cases.

$\sigma(s)$	$\tau(s)$	$\varrho(s)$	α, β	(a, b)
1	$\alpha s + \beta$	$\mathrm{e}^{\alpha s^2/2+\beta s}$	$\alpha < 0$	$(-\infty, \infty)$
s	$\alpha s + \beta$	$s^{\beta-1}\mathrm{e}^{\alpha s}$	$\alpha < 0,\ \beta > 0$	$(0, \infty)$
$1 - s^2$	$\alpha s + \beta$	$(1+s)^{-(\alpha-\beta)/2-1}(1-s)^{-(\alpha+\beta)/2-1}$	$\alpha < \beta < -\alpha$	$(-1, 1)$
$s^2 - 1$	$\alpha s + \beta$	$(s+1)^{(\alpha-\beta)/2-1}(s-1)^{(\alpha+\beta)/2-1}$	$-\beta < \alpha < 0$	$(1, \infty)$
s^2	$\alpha s + \beta$	$s^{\alpha-2}\mathrm{e}^{-\beta/s}$	$\alpha < 0,\ \beta > 0$	$(0, \infty)$
$s^2 + 1$	$\alpha s + \beta$	$(1+s^2)^{\alpha/2-1}\mathrm{e}^{\beta\arctan s}$	$\alpha < 0$	$(-\infty, \infty)$

Our main purpose is to present a unified view on the systems of orthogonal polynomials defined by equation (1) in the cases presented in table 1. We analyse the associated special functions, the corresponding raising/lowering operators, and present some applications to quantum mechanics. Some of our results and proofs are extended versions of some results and proofs presented in [5]. The general form of the raising/lowering operators has been previously obtained by Jafarizadeh and Fakhri by using a different method [4]. Our results concerning the relation with classical polynomials obtained in a direct way are in agreement with those based on recursion relations presented in [1, 2].

2. Polynomials of hypergeometric type

It is well-known [5] that for $\lambda = \lambda_l$, where

$$\lambda_l = -\frac{\sigma''(s)}{2}l(l-1) - \tau'(s)l \qquad l \in \mathbb{N} \tag{4}$$

the equation (1) admits a polynomial solution $\Phi_l = \Phi_l^{(\alpha,\beta)}$ of at most l degree

$$\sigma(s)\Phi_l'' + \tau(s)\Phi_l' + \lambda_l\Phi_l = 0. \tag{5}$$

If the degree of the polynomial Φ_l is l then it satisfies the Rodrigues formula

$$\Phi_l(s) = \frac{B_l}{\varrho(s)}[\sigma^l(s)\varrho(s)]^{(l)} \tag{6}$$

where B_l is a constant. We do not impose any normalizing condition. Our polynomials Φ_l are defined only up to a multiplicative constant. One can remark that

$$\lim_{s\to a}\sigma(s)\varrho(s)s^\gamma = \lim_{s\to b}\sigma(s)\varrho(s)s^\gamma = 0 \qquad \text{for} \quad \gamma \in [0,\infty) \tag{7}$$

in the case $\sigma(s) \in \{1,\ s,\ 1-s^2\}$, and

$$\lim_{s\to a}\sigma(s)\varrho(s)s^\gamma = \lim_{s\to b}\sigma(s)\varrho(s)s^\gamma = 0 \qquad \text{for} \quad \gamma \in [0,-\alpha) \tag{8}$$

in the case $\sigma(s) \in \{s^2-1,\ s^2,\ s^2+1\}$. Let

$$\nu = \begin{cases} \infty & \text{for} \quad \sigma(s) \in \{1,\ s,\ 1-s^2\} \\ \frac{1-\alpha}{2} & \text{for} \quad \sigma(s) \in \{s^2-1,\ s^2,\ s^2+1\}. \end{cases} \tag{9}$$

Theorem 1.
a) $\{\Phi_l \mid l < \nu\}$ *is a system of polynomials orthogonal with weight function* $\varrho(s)$ *in* (a,b).

b) Φ_l *is a polynomial of degree* l *for any* $l < \nu$.

c) The function $\Phi_l(s)\sqrt{\varrho(s)}$ *is square integrable on* (a,b) *for any* $l < \nu$.

d) A three term recurrence relation

$$s\Phi_l(s) = \alpha_l\Phi_{l+1}(s) + \beta_l\Phi_l(s) + \gamma_l\Phi_{l-1}(s)$$

is satisfied for $1 < l+1 < \nu$.

e) The zeros of Φ_l *are simple and lie in the interval* (a,b), *for any* $l < \nu$.

Proof. a) Let $l,\ k \in \mathbb{N}$ with $0 \le l < k < \nu$. From the relations

$$[\sigma(s)\varrho(s)\Phi_l']' + \lambda_l\varrho(s)\Phi_l = 0 \qquad [\sigma(s)\varrho(s)\Phi_k']' + \lambda_k\varrho(s)\Phi_k = 0$$

we get

$$(\lambda_l - \lambda_k)\Phi_l(s)\Phi_k(s)\varrho(s) = \frac{d}{ds}\{\sigma(s)\varrho(s)[\Phi_l(s)\Phi_k'(s) - \Phi_k(s)\Phi_l'(s)]\}.$$

Since the Wronskian $W[\Phi_l(s),\Phi_k(s)] = \Phi_l(s)\Phi_k'(s) - \Phi_k(s)\Phi_l'(s)$ is a polynomial of at most $l+k-1$ degree, from (7) and (8) it follows that

$$(\lambda_l - \lambda_k)\int_a^b \Phi_l(s)\Phi_k(s)\varrho(s)ds$$

$$= \lim_{s\to b} \sigma(s)\varrho(s)W[\Phi_l(s), \Phi_k(s)] - \lim_{s\to a} \sigma(s)\varrho(s)W[\Phi_l(s), \Phi_k(s)] = 0.$$

We have $\lambda_l - \lambda_k \neq 0$ since the function $l \mapsto \lambda_l$ is strictly increasing on $\{l \in \mathbb{N} \mid l < \nu\}$.
b) Each Φ_l is a polynomial of at most l degree and the polynomials $\{\Phi_l \mid 0 \leq l < \nu\}$ are linearly independent. This is possible only if Φ_l is a polynomial of degree l for any $l < \nu$.
c) In the case $\sigma(s) \in \{s^2 - 1,\ s^2,\ s^2 + 1\}$ we have $l < (1 - \alpha)/2$, that is, $2l - 1 < -\alpha$. Therefore there exists $\varepsilon > 0$ such that $1 + \varepsilon + 2l - 2 = 2l - 1 + \varepsilon < -\alpha$, and hence

$$\lim_{s\to\infty} s^{1+\varepsilon}|\Phi_l(s)|^2\varrho(s) = \lim_{s\to\infty} \frac{s^{1+\varepsilon}(\Phi_l(s))^2}{\sigma(s)}\sigma(s)\varrho(s) = 0.$$

The convergence of the integral

$$\int_a^b |\Phi_l(s)|^2\varrho(s)ds$$

follows from the convergence of the integral $\int_1^\infty s^{-1-\varepsilon}ds$. In the case $\sigma(s) = 1 - s^2$, from $\alpha < \beta < -\alpha$ we get $-(\alpha - \beta)/2 - 1 > -1$ and $-(\alpha + \beta/2 - 1 > -1$.
d,e) See [5].

Theorem 2. *Up to a multiplicative constant*

$$\Phi_l^{(\alpha,\beta)}(s) = \begin{cases} H_l\left(\sqrt{\frac{-\alpha}{2}}\, s - \frac{\beta}{\sqrt{-2\alpha}}\right) & \text{in the case} \quad \sigma(s) = 1 \\ L_l^{\beta-1}(-\alpha s) & \text{in the case} \quad \sigma(s) = s \\ P_l^{(-(\alpha+\beta)/2-1,\ (-\alpha+\beta)/2-1)}(s) & \text{in the case} \quad \sigma(s) = 1 - s^2 \\ P_l^{((\alpha-\beta)/2-1,\ (\alpha+\beta)/2-1)}(-s) & \text{in the case} \quad \sigma(s) = s^2 - 1 \\ \left(\frac{s}{\beta}\right)^l L_l^{1-\alpha-2l}\left(\frac{\beta}{s}\right) & \text{in the case} \quad \sigma(s) = s^2 \\ \mathrm{i}^l P_l^{((\alpha+\mathrm{i}\beta)/2-1,\ (\alpha-\mathrm{i}\beta)/2-1)}(\mathrm{i}s) & \text{in the case} \quad \sigma(s) = s^2 + 1 \end{cases} \tag{10}$$

where H_n, L_n^p and $P_n^{(p,q)}$ are the Hermite, Laguerre and Jacobi polynomials, respectively.

Proof. In the case $\sigma(s) = s^2$ the function $\Phi_l^{(\alpha,\beta)}(s)$ satisfies the equation

$$s^2y'' + (\alpha s + \beta)y' + [-l(l-1) - \alpha l]y = 0.$$

If we denote $t = \beta/s$ then the polynomial $u(t) = t^l y(\beta/t)$ satisfies the equation

$$tu'' + (-\alpha + 2 - 2l - t)u' + lu = 0$$

that is, the equation whose polynomial solution is $L_l^{1-\alpha-2l}(s)$ (up to a multiplicative constant). In a similar way one can analyse the other cases.

3. Special functions of hypergeometric type

Let $l \in \mathbb{N}$, $l < \nu$, and let $m \in \{0, 1, ..., l\}$. By differentiating the equation (5) m times we obtain the equation satisfied by the polynomials $\varphi_{l,m} = \Phi_l^{(m)}$, namely

$$\sigma(s)\varphi''_{l,m} + [\tau(s) + m\sigma'(s)]\varphi'_{l,m} + (\lambda_l - \lambda_m)\varphi_{l,m} = 0. \tag{11}$$

This is an equation of hypergeometric type, and we can write it in the self-adjoint form

$$[\sigma(s)\varrho_m(s)\varphi'_{l,m}]' + (\lambda_l - \lambda_m)\varrho_m(s)\varphi_{l,m} = 0 \tag{12}$$

by using the function $\varrho_m(s) = \sigma^m(s)\varrho(s)$.

The functions

$$\Phi_{l,m}(s) = \kappa^m(s)\Phi_l^{(m)}(s) \tag{13}$$

where $l \in \mathbb{N}$, $l < \nu$, $m \in \{0, 1, ..., l\}$ and $\kappa(s) = \sqrt{\sigma(s)}$, are called the *associated special functions*. The equation (11) multiplied by $\kappa^m(s)$ can be written as

$$H_m\Phi_{l,m} = \lambda_l\Phi_{l,m} \tag{14}$$

where H_m is the differential operator

$$\begin{aligned} H_m = & -\sigma(s)\frac{d^2}{ds^2} - \tau(s)\frac{d}{ds} + \frac{m(m-2)}{4}\frac{\sigma'^2(s)}{\sigma(s)} \\ & + \frac{m\tau(s)}{2}\frac{\sigma'(s)}{\sigma(s)} - \frac{1}{2}m(m-2)\sigma''(s) - m\tau'(s). \end{aligned} \tag{15}$$

Theorem 3. *a) For each $m < \nu$, the functions $\Phi_{l,m}$ with $m \leq l < \nu$ are orthogonal with weight function $\varrho(s)$ in (a, b).*

b) $\Phi_{l,m}(s)\sqrt{\varrho(s)}$ is square integrable on (a, b) for $0 \leq m \leq l < \nu$.

c) The three term recurrence relation

$$\Phi_{l,m+1}(s) + \left(\frac{\tau(s)}{\kappa(s)} + 2(m-1)\kappa'(s)\right)\Phi_{l,m}(s) + (\lambda_l - \lambda_{m-1})\Phi_{l,m-1}(s) = 0 \tag{16}$$

is satisfied for any $l < \nu$ and any $m \in \{1, 2, ..., l-1\}$. In addition, we have

$$\left(\frac{\tau(s)}{\kappa(s)} + 2(l-1)\kappa'(s)\right)\Phi_{l,l}(s) + (\lambda_l - \lambda_{l-1})\Phi_{l,l-1}(s) = 0. \tag{17}$$

Proof. a) In the case $\sigma(s) \in \{s^2 - 1,\ s^2,\ s^2 + 1\}$, for each $k < -\alpha - 2m$ we have

$$\lim_{s\to a} \sigma(s)\varrho_m(s)s^k = \lim_{s\to b} \sigma(s)\varrho_m(s)s^k = 0. \tag{18}$$

Since the equation (11) is of hypergeometric type, $\varphi_{l,m}$ is a polynomial of degree $l - m$, and $\varphi_{k,m}$ is a polynomial of degree $k - m$, from theorem 1 applied to (11) it follows that

$$\int_a^b \varphi_{l,m}(s)\varphi_{k,m}(s)\varrho_m(s)ds = 0$$

for $l \neq k$ with $(l-m)+(k-m) < -\alpha-2m$, that is, $l+k < -\alpha$. But $\varrho_m(s) = \sigma^m(s)\varrho(s)$, and hence

$$\int_a^b \varphi_{l,m}(s)\varphi_{k,m}(s)\varrho_m(s)ds = \int_a^b \Phi_{l,m}(s)\Phi_{k,m}(s)\varrho(s)ds.$$

b) In the case $\sigma(s) \in \{s^2 - 1,\ s^2,\ s^2 + 1\}$, for $\varepsilon > 0$ chosen such that $2\nu + \varepsilon < 1 - \alpha$ we have $1 + \varepsilon + 2(m-1) + 2(l-m) \leq 1 + \varepsilon + 2\nu - 2 < -\alpha$, whence

$$\lim_{s\to\infty} s^{1+\varepsilon}(\Phi_{l,m}(s))^2\varrho(s) = \lim_{s\to\infty} s^{1+\varepsilon}\sigma^{m-1}(s)(\Phi_l^{(m)}(s))^2\sigma(s)\varrho(s) = 0.$$

The convergence of the integral

$$\int_a^b |\Phi_{l,m}(s)|^2 \varrho(s)ds$$

follows from the convergence of the integral $\int_1^\infty s^{-1-\varepsilon}ds$.
c) By differentiating (5) $m-1$ times we obtain

$$\sigma(s)\Phi_l^{(m+1)}(s) + (m-1)\sigma'(s)\Phi_l^{(m)}(s) + \frac{(m-1)(m-2)}{2}\sigma''(s)\Phi_l^{(m-1)}(s)$$

$$+\tau(s)\Phi_l^{(m)} + (m-1)\tau'(s)\Phi_l^{(m-1)}(s) + \lambda_l\Phi_l^{(m-1)}(s) = 0.$$

If we multiply this relation by $\kappa^{m-1}(s)$ then we get (16) for $m \in \{1, 2, ..., l-1\}$, and (17) for $m = l$.

4. Raising and lowering operators

For any $l \in \mathbb{N}$, $l < \nu$ and any $m \in \{0, 1, ..., l-1\}$, by differentiating (13), we obtain

$$\Phi'_{l,m}(s) = m\kappa^{m-1}(s)\kappa'(s)\Phi_l^{(m)} + \kappa^m(s)\Phi_l^{(m+1)}(s)$$

that is, the relation

$$\Phi'_{l,m}(s) = m\frac{\kappa'(s)}{\kappa(s)}\Phi_{l,m}(s) + \frac{1}{\kappa(s)}\Phi_{l,m+1}(s)$$

which can be written as

$$\left(\kappa(s)\frac{d}{ds} - m\kappa'(s)\right)\Phi_{l,m}(s) = \Phi_{l,m+1}(s). \tag{19}$$

If $m \in \{1, 2, ..., l-1\}$ then by substituting (19) into (16) we get

$$\left(\kappa(s)\frac{d}{ds} + \frac{\tau(s)}{\kappa(s)} + (m-2)\kappa'(s)\right)\Phi_{l,m}(s) + (\lambda_l - \lambda_{m-1})\Phi_{l,m-1}(s) = 0$$

that is,

$$\left(-\kappa(s)\frac{d}{ds} - \frac{\tau(s)}{\kappa(s)} - (m-1)\kappa'(s)\right)\Phi_{l,m+1}(s) = (\lambda_l - \lambda_m)\Phi_{l,m}(s). \tag{20}$$

for all $m \in \{0, 1, ..., l-2\}$. From (17) it follows that this relation is also satisfied for $m = l-1$.

The relations (19) and (20) suggest us to consider the first order differential operators

$$A_m = \kappa(s)\frac{d}{ds} - m\kappa'(s) \qquad A_m^+ = -\kappa(s)\frac{d}{ds} - \frac{\tau(s)}{\kappa(s)} - (m-1)\kappa'(s) \tag{21}$$

for $m+1 < \nu$.

Theorem 4. *We have*

a) $\quad A_m\Phi_{l,m} = \Phi_{l,m+1} \qquad A_m^+\Phi_{l,m+1} = (\lambda_l - \lambda_m)\Phi_{l,m} \qquad$ *for* $\quad 0 \leq m < l < \nu.$ (22)

$$b)\quad \Phi_{l,m}=\frac{A_m^+}{\lambda_l-\lambda_m}\frac{A_{m+1}^+}{\lambda_l-\lambda_{m+1}}\cdots\frac{A_{l-2}^+}{\lambda_l-\lambda_{l-2}}\frac{A_{l-1}^+}{\lambda_l-\lambda_{l-1}}\Phi_{l,l}\qquad for\quad 0\leq m<l<\nu. \tag{23}$$

$$c)\quad ||\Phi_{l,m+1}||=\sqrt{\lambda_l-\lambda_m}\,||\Phi_{l,m}||\qquad for\quad 0\leq m<l<\nu. \tag{24}$$

$$d)\quad H_m-\lambda_m=A_m^+A_m\qquad H_{m+1}-\lambda_m=A_mA_m^+\qquad for\quad m+1<\nu \tag{25}$$

$$e)\quad H_mA_m^+=A_m^+H_{m+1}\qquad A_mH_m=H_{m+1}A_m\qquad for\quad m+1<\nu. \tag{26}$$

Proof. a) These relations coincide to (19) and (20), respectively.
b) Since the function $k\mapsto\lambda_k$ is strictly increasing on $\{k\in\mathbb{N}\mid k<\nu\}$ we have $\lambda_l-\lambda_k\neq 0$ for $k\neq l$. The formula (23) follows from (22).
c) Since $\sigma^m(s)\Phi_l^{(m)}(s)\Phi_k^{(m+1)}(s)$ is a polynomial of degree $l+k-1$, from (7) and (8) we get

$$\begin{aligned}\langle A_m\Phi_{l,m},\Phi_{k,m+1}\rangle&=\int_a^b[\kappa(s)\Phi'_{l,m}(s)-m\kappa'(s)\Phi_{l,m}(s)]\Phi_{k,m+1}(s)\varrho(s)ds\\&=\kappa(s)\Phi_{l,m}(s)\Phi_{k,m+1}(s)\varrho(s)|_a^b-\int_a^b\Phi_{l,m}(s)[\kappa(s)\Phi'_{k,m+1}(s)\varrho(s)\\&+\kappa(s)\Phi_{k,m+1}(s)\varrho'(s)+(m+1)\kappa'(s)\Phi_{k,m+1}(s)\varrho(s)]ds\\&=\sigma(s)\varrho(s)\sigma^m(s)\Phi_l^{(m)}(s)\Phi_k^{(m+1)}(s)|_a^b+\int_a^b\Phi_{l,m}(s)(A_m^+\Phi_{k,m+1})(s)\varrho(s)ds\\&=\langle\Phi_{l,m},A_m^+\Phi_{k,m+1}\rangle\end{aligned}$$

whence

$$||\Phi_{l,m+1}||^2=\langle\Phi_{l,m+1},\Phi_{l,m+1}\rangle=\langle A_m\Phi_{l,m},\Phi_{l,m+1}\rangle=\langle\Phi_{l,m},A_m^+\Phi_{l,m+1}\rangle=(\lambda_l-\lambda_m)||\Phi_{l,m}||^2.$$

d,e) These relations can be proved by direct computation.

5. Application to Schrödinger type operators

If we use in equation (14) a change of variable $(a',b')\longrightarrow(a,b):x\mapsto s(x)$ such that $ds/dx=\kappa(s(x))$ or $ds/dx=-\kappa(s(x))$ and define the new functions

$$\Psi_{l,m}(x)=\sqrt{\kappa(s(x))\,\varrho(s(x))}\,\Phi_{l,m}(s(x)) \tag{27}$$

then we get an equation of Schrödinger type [3]

$$-\frac{d^2}{dx^2}\Psi_{l,m}(x)+V_m(x)\Psi_{l,m}(x)=\lambda_l\Psi_{l,m}(x). \tag{28}$$

Since

$$\int_{a'}^{b'}|\Psi_{l,m}(x)|^2dx=\int_{a'}^{b'}|\Phi_{l,m}(s(x))|^2\varrho(s(x))\frac{d}{dx}s(x)dx=\int_a^b|\Phi_{l,m}(s)|^2\varrho(s)ds$$

$$\int_{a'}^{b'}\Psi_{l,m}(x)\Psi_{k,m}(x)dx=\int_a^b\Phi_{l,m}(s)\Phi_{k,m}(s)\varrho(s)ds$$

the functions $\Psi_{l,m}(x)$ with $0 \leq m \leq l < \nu$ are square integrable on (a', b') and orthogonal.

If $ds/dx = \kappa(s(x))$ then the operators corresponding to A_m and A_m^+ are

$$\begin{array}{l} \mathcal{A}_m = [\kappa(s)\varrho(s)]^{1/2} A_m [\kappa(s)\varrho(s)]^{-1/2}|_{s=s(x)} = \frac{d}{dx} + W_m(x) \\ \mathcal{A}_m^+ = [\kappa(s)\varrho(s)]^{1/2} A_m^+ [\kappa(s)\varrho(s)]^{-1/2}|_{s=s(x)} = -\frac{d}{dx} + W_m(x) \end{array} \tag{29}$$

where the *superpotential* $W_m(x)$ is given by the formula [4]

$$W_m(x) = -\frac{\tau(s(x))}{2\kappa(s(x))} - \frac{2m-1}{2\kappa(s(x))} \frac{d}{dx}\kappa(s(x))\,. \tag{30}$$

From (22) and (25) we get

$$\mathcal{A}_m \Psi_{l,m}(x) = \Psi_{l,m+1}(x) \qquad \mathcal{A}_m^+ \Psi_{l,m+1}(x) = (\lambda_l - \lambda_m)\Psi_{l,m}(x) \tag{31}$$

and

$$(\mathcal{A}_m^+ \mathcal{A}_m + \lambda_m)\Psi_{l,m} = \lambda_l \Psi_{l,m} \qquad (\mathcal{A}_m \mathcal{A}_m^+ + \lambda_m)\Psi_{l,m+1} = \lambda_l \Psi_{l,m+1} \tag{32}$$

whence

$$-\frac{d^2}{dx^2} + V_m(x) - \lambda_m = \mathcal{A}_m^+ \mathcal{A}_m \qquad -\frac{d^2}{dx^2} + V_{m+1}(x) - \lambda_m = \mathcal{A}_m \mathcal{A}_m^+ \tag{33}$$

and

$$V_m(x) - \lambda_m = W_m^2(x) - \dot{W}_m(x) \qquad V_{m+1}(x) - \lambda_m = W_m^2(x) + \dot{W}_m(x) \tag{34}$$

where the dot sign means derivative with respect to x.

Since $\mathcal{A}_m \Psi_{m,m} = 0$, from (29) and (33) we get

$$\dot{\Psi}_{m,m} + W_m(x)\Psi_{m,m} = 0 \qquad -\ddot{\Psi}_{m,m} + (V_m(x) - \lambda_m)\Psi_{m,m} = 0 \tag{35}$$

whence

$$W_m(x) = -\frac{\dot{\Psi}_{m,m}(x)}{\Psi_{m,m}(x)} \qquad V_m(x) = \frac{\ddot{\Psi}_{m,m}(x)}{\Psi_{m,m}(x)} + \lambda_m. \tag{36}$$

For each $l \in \{0, 1, ..., \nu\}$ and each $m \in \{0, 1, ..., l-1\}$ we have

$$\Psi_{l,m}(x) = \frac{\mathcal{A}_m^+}{\lambda_l - \lambda_m} \frac{\mathcal{A}_{m+1}^+}{\lambda_l - \lambda_{m+1}} ... \frac{\mathcal{A}_{l-2}^+}{\lambda_l - \lambda_{l-2}} \frac{\mathcal{A}_{l-1}^+}{\lambda_l - \lambda_{l-1}} \Psi_{l,l}(x). \tag{37}$$

If we choose the change of variable $s = s(x)$ such that $ds/dx = -\kappa(s(x))$, then the formulae (29), (30), (34) and (36) become

$$\mathcal{A}_m = -\frac{d}{dx} + W_m(x) \qquad \mathcal{A}_m^+ = \frac{d}{dx} + W_m(x) \tag{38}$$

$$W_m(x) = -\frac{\tau(s(x))}{2\kappa(s(x))} + \frac{2m-1}{2\kappa(s(x))} \frac{d}{dx}\kappa(s(x)) \tag{39}$$

$$V_m(x) - \lambda_m = W_m^2(x) + \dot{W}_m(x) \qquad V_{m+1}(x) - \lambda_m = W_m^2(x) - \dot{W}_m(x) \tag{40}$$

$$W_m(x) = \frac{\dot{\Psi}_{m,m}(x)}{\Psi_{m,m}(x)} \qquad V_m(x) = \frac{\ddot{\Psi}_{m,m}(x)}{\Psi_{m,m}(x)} + \lambda_m \tag{41}$$

respectively.

Examples. Let $\alpha_m = (1-\alpha-2m)/2,\ \ \alpha'_m = (-1-\alpha+2m)/2$ and $\delta = -\beta/2$.

1. In the case $\sigma(s) = 1-s^2,\ \ \tau(s) = \alpha s+\beta,\ \ s(x) = \cos x$ we get

$$W_m(x) = \alpha'_m \ \mathrm{cotan}\, x + \delta\ \mathrm{cosec}\, x = \tfrac{\alpha'_m+\beta}{2}\ \mathrm{cotan}\, \tfrac{x}{2} - \tfrac{\alpha'_m-\beta}{2} \tan \tfrac{x}{2}$$

$$V_m(x) = \left({\alpha'_m}^2 - \alpha'_m + \delta^2\right)\ \mathrm{cosec}^2 x + (2\alpha_m - 1)\delta\ \mathrm{cotan}\, x\ \mathrm{cosec}\, x - {\alpha'_m}^2 + \lambda_m$$

$$(Poschl - Teller\ type\ potential).$$

2. In the case $\sigma(s) = s^2-1,\ \ \tau(s) = \alpha s+\beta,\ \ s(x) = \cosh x$ we get

$$W_m(x) = \alpha_m\ \mathrm{cotanh}\ x + \delta\ \mathrm{cosech}\, x$$

$$V_m(x) = (\alpha_m^2 + \alpha_m + \delta^2)\ \mathrm{cosech}^2 x + (2\alpha_m + 1)\delta\ \mathrm{cotanh}\ x\ \mathrm{cosech}\, x + \alpha_m^2 + \lambda_m$$

$$(generalized\ Poschl - Teller\ potential).$$

3. In the case $\sigma(s) = s^2,\ \ \tau(s) = \alpha s+\beta,\ \ s(x) = \mathrm{e}^x$ we get

$$W_m(x) = \alpha_m + \delta \mathrm{e}^{-x}$$

$$V_m(x) = \delta^2 \mathrm{e}^{-2x} + (2\alpha_m + 1)\delta \mathrm{e}^{-x} + \alpha_m^2 + \lambda_m$$

$$(Morse\ type\ potential).$$

4. In the case $\sigma(s) = s^2+1,\ \ \tau(s) = \alpha s+\beta,\ \ s(x) = \sinh x$ we get

$$W_m(x) = \alpha_m \tanh x + \delta\ \mathrm{sech}\, x$$

$$V_m(x) = (-\alpha_m^2 - \alpha_m + \delta^2)\ \mathrm{sech}^2 x + (2\alpha_m + 1)\delta \tanh x\ \mathrm{sech}\, x + \alpha_m^2 + \lambda_m$$

$$(Scarf\ hyperbolic\ type\ potential).$$

6. Concluding remarks

The equation (1) defines an infinite system of orthogonal polynomials in the first three cases presented in table 1, and a finite system of orthogonal polynomials in the last three cases. Despite the fact that all these orthogonal polynomials can be expressed in terms of the classical ones (theorem 2), we think that it is worth considering them. They are directly related to the bound-state eigenfunctions of some important Schrödinger equations and allow us to analyse these equations together, in a unified formalism. The relation with classical polynomials (10) has not a very simple form in all the cases. More than that, in certain cases we have to consider the classical polynomials outside the interval where they are orthogonal or for complex values of parameters. Generally, the properties of the functions considered in this paper (orthogonality, square integrability, recursion relations, raising/lowering operators) can not be obtained in a simple way from those concerning the classical polynomials.

Acknowledgments

This research was supported by the grant CERES no. 24/2002.

References

[1] Cooper F, Khare A and Sukhatme U 1995 Supersymmetry and quantum mechanics *Phys. Rep.* **251** 267–385

[2] Dabrowska J W, Khare A and Sukhatme 1988 Explicit wavefunctions for shape-invariant potentials by operator techniques J. Phys. A: Math. Gen. **21** L195-L200

[3] Infeld L and Hull T E 1951 The factorization method *Rev. Mod. Phys.* **23** 21–68

[4] Jafarizadeh M A and Fakhri H 1998 Parasupersymmetry and shape invariance in differential equations of mathematical physics and quantum mechanics *Ann. Phys., NY* **262** 260–76

[5] Nikiforov A F, Suslov S K and Uvarov V B 1991 *Classical Orthogonal Polynomials of a Discrete Variable* (Berlin: Springer)